

Possible Glassiness in a Periodic Long-Range Josephson Array

P. Chandra

NEC Research Institute, 4 Independence Way, Princeton NJ 08540

L. B. Ioffe

*Department of Physics, Rutgers University, Piscataway, NJ 08855
and Landau Institute for Theoretical Physics, Moscow*

D. Sherrington

Department of Physics, Oxford University, 1 Keble Rd, Oxford, OX1 3NP, UK

We present an analytic study of a periodic Josephson array with long-range interactions in a transverse magnetic field. We find that this system exhibits a first-order transition into a phase characterized by an extensive number of states separated by barriers that scale with the system size; the associated discontinuity is small in the limit of weak applied field, thus permitting an explicit analysis in this regime.

The essential features underlying the physics of glass formation are still unclear. In particular the possibility of glassiness in the absence of disorder is a question that has stimulated much recent activity [1]. In this paper we present and study a periodic Josephson array with long-range interactions frustrated by a transverse magnetic field (H). We show that it displays a first-order transition into a low-temperature phase characterized by an extensive number of states; these metastable solutions are separated by barriers that scale with the system size. The discontinuity at the transition is small in the limit of weak applied field, so that an explicit analytic treatment is possible in this regime.




Fig 1. A schematic of the array discussed in the text; the solid (shaded) lines are the horizontal (vertical) superconducting wires that are coupled by Josephson junctions to the vertical (horizontal) wires at each node.

More specifically, the proposed array is a stack of two mutually perpendicular sets of N parallel wires with Josephson junctions at each node (Figure 1) that is placed in an external transverse field. The classical thermodynamic variables of this system are the superconducting phases associated with each wire. Here we shall assume that the Josephson couplings are sufficiently small so that the induced fields are negligible in comparison with H . We can therefore describe the array by the Hamiltonian

$$\mathcal{H} = -\text{Re} \sum_{j,k} S_j J_{jk} S_k = - \sum_{m,n} s_m^* \mathcal{J}_{mn} s_n \quad (1)$$

where $\mathcal{J}_{mn}$ is the coupling matrix

$$\hat{\mathcal{J}} = \begin{pmatrix} 0 & \hat{J} \\ \hat{J}^\dagger & 0 \end{pmatrix} \quad (2)$$

with $J_{jk} = \frac{J_0}{\sqrt{N}} \exp(2\pi i \alpha j k / N)$ and $S_j(s_m)$ is a $N(2N)$ -element vector whose components are $\exp(i\phi_m)$, where ϕ_m are the superconducting phases of the wires. Here we have introduced the flux per unit strip, $\alpha = N H l^2 / \phi_0$, where l is the inter-node spacing, ϕ_0 is the flux quantum, and the normalization has been chosen so that $T_c \sim J_0$.

Because every horizontal (vertical) wire is linked to every vertical (horizontal) wire, the number of nearest neighbors in this model $z = N$; we can therefore study it with a mean-field approach. For the same reason the free energy barriers separating its low T solutions scale with N , in marked contrast to the situation in a conventional 2D array [2]; thus each of these solutions results in a metastable state. If the field is such that the number of solutions is extensive (i.e. the phase-ordering unit cell scales with the system size) this array displays glassy behavior in the absence of disorder.

A similar long-range network with areal disorder was studied previously [3]; for $\alpha \gg 1/N$ the system undergoes a spin glass transition which was mapped onto the Sherrington-Kirkpatrick model for the case $\alpha \gg 1$ for all T . Physically this behavior occurs because the phase differences associated with the couplings, J_{jk} , acquire random values and fill the interval $(0, 2\pi)$ uniformly for $\alpha \gg 1/N$. In the absence of randomness these phases still cover the interval for $\alpha \gg 1/N$, so that one expects that glassy behavior survive [4]. Naturally, there will be special values of the field for which the number of solutions is finite. However there is no such commensurability for $1/N \ll \alpha \leq 1$; e.g. the unit cell corresponding to $\alpha = 1$ consists of all the wires traversing a strip and thus has sides of order $O(N)l$.

The cumulants of the coupling constants display a more quantitative similarity between the random and the periodic arrays. In both cases

$$\sum_{j_1, j_2, \dots, j_n} J_{i_1 j_1} \dots J_{j_n i_1} = J_0^{2n} \left(\frac{1}{\alpha} \right)^{n-1} \quad (3)$$

which can be understood physically since only horizontal (vertical) wires separated by $d < l/\alpha$ contribute coherently to this sum independent of disorder.

We now begin a more quantitative analysis of this periodic array using a modified Thouless-Anderson-Palmer approach [5]. Because the interactions are long-range, we can integrate out the thermal fluctuations and define a “site magnetization” $m_i = \langle z_i \rangle_T$. The free energy is

$$\begin{aligned} F\{m_k\} = & - \sum_{kj} m_k^* \mathcal{J}_{kj} m_j - T \sum_k S(m_k) + F_o\{m_k\} \\ & - \sum_k (h_k m_k^* + h_k^* m_k) \end{aligned} \quad (4)$$

$$S(m) = S_0 - |m|^2 - \frac{1}{4}|m|^4 + O(|m|^6)$$

where $\mathcal{J}_{kj}$ is defined in (2). Here we have introduced h_i , a field conjugate to m_i , and the single site entropy $S(m)$; $F_o\{m_k\}$ is the Onsager reaction term. At high temperatures $F_o\{m_k\} \sim 1/T$ and thus is small; furthermore, as we show below, the transition occurs at $T_c \approx J_0/\sqrt{\alpha}$ (as in the disordered case) so that we expect the feedback effects to be perturbative for small α ($1/N \ll \alpha \ll 1$). Thus we shall ignore $F_o\{m_k\}$ in our initial study of $F\{m_k\}$, but will consider it later.

In the high temperature phase $m_i \equiv 0$. This “paramagnetic” state becomes unstable when the quadratic part of $F\{m_k\}$ acquires negative eigenvalues, i.e. at $T \leq \lambda_{max}$ where λ_{max} is the largest eigenvalue of $\hat{\mathcal{J}}$. Due to the structure of $\hat{\mathcal{J}}$ (see (2)), any eigenvalue Λ of $\hat{\mathcal{J}}\hat{\mathcal{J}}^\dagger$ results in two eigenvalues $\lambda = \pm\sqrt{\Lambda}$ of $\hat{\mathcal{J}}$, a property we exploit for convenience. In the limit $N \rightarrow \infty$

$$\left(\hat{\mathcal{J}}\hat{\mathcal{J}}^\dagger \right)_{jk} = J_0^2 e^{i\pi\alpha(j-k)} \frac{\sin(\pi\alpha(j-k))}{\pi\alpha(j-k)} \quad (5)$$

which can be diagonalized by a Fourier transformation to yield

$$\left(\hat{\mathcal{J}}\hat{\mathcal{J}}^\dagger \right)_p = \frac{J_0^2}{\alpha} \theta(p) \theta(2\pi\alpha - p). \quad (6)$$

Therefore the largest eigenvalue of $\hat{\mathcal{J}}$ is $\lambda_{max} = J_0/\sqrt{\alpha}$; the corresponding eigenfunctions associated with one set of parallel wires are plane waves with momenta in the interval $0 \leq p \leq 2\pi\alpha$ so that the degeneracy of this eigenvalue is αN [6].

Therefore, in the absence of the feedback effects, the transition occurs at $T_{c0} = J_0/\sqrt{\alpha}$ into a phase that is characterized by a linear combination of the αN states described above. Thus the number of possible metastable configurations just below the transition should be extensive, indicating the onset of glassiness [7]. This should be contrasted with the situation in the Sherrington-Kirkpatrick model where, in the thermodynamic limit, the highest eigenvalue of the exchange matrix is non-degenerate and the metastability measure of the order parameter (plateau-onset x_0 in Parisi function $q(x)$) grows continuously beneath the transition. More generally, in disordered spin glasses a sudden appearance of extensive metastability implies a first-order jump in the order parameter [8]. Unfortunately in the periodic case we have not yet determined the analogous order parameter; however we show below that the Onsager feedback term drives the transition weakly first-order, consistent with our expectations.

We use the locator expansion [9] to determine the leading order contributions in m_i to $F_o\{m_k\}$, it is based on the expression for susceptibility

$$\hat{\chi} = \frac{1}{\hat{A} - \hat{\mathcal{J}}} \quad (7)$$

where $\hat{A}$ is a diagonal “locator” matrix. This form of susceptibility is a consequence of large z ; to prove it one can use the high temperature expansion and show that all diagrams renormalizing the $\hat{\mathcal{J}}$ term are small in $1/z$. Once we have solved for $\hat{A}$, we can reconstruct the free energy from

$$\frac{\partial^2 F}{\partial m_k^* \partial m_j} = (\hat{\chi}^{-1})_{kj} \quad (8)$$

that gives us the free energy which contains both the entropy term $\sum T S(m_i)$ and the Onsager term $F_o\{m_i\}$. Although it is possible, in principle, to determine the elements of $\hat{A}$ *ab initio* it is easier to do so using the identity $\chi_{jj} \equiv \frac{1-|m_j|^2}{T}$ which follows from the fluctuation-dissipation theorem. We expand the expression for the susceptibility (7) and, due to the structure of $\hat{\mathcal{J}}$, find that only terms even in $\hat{\mathcal{J}}$ contribute. Collecting and resumming these terms, we obtain

$$\left(\frac{1}{\hat{A} - \hat{J}\hat{A}^{-1}\hat{J}^\dagger} \right)_{jj} = \frac{1 - |m_j|^2}{T}. \quad (9)$$

We first consider the leading-order term (setting $m_j = 0$) which will yield the quadratic part of $F\{m_j\}$. To this order $\hat{A}$ is independent of the site label ($\hat{A} = A\hat{1}$), so it is convenient to solve this equation by Fourier transformation and, using (5), we obtain

$$\frac{\alpha}{A - \frac{J_0^2}{A\alpha}} + \frac{1 - \alpha}{A} = \frac{1}{T} \quad (10)$$

The zeroth-order term in α yields $A_0 = T$, consistent with our assumption above that the reaction term is small in α .




Fig 2. Plot of the locator $A(T)$ obtained as a numerical solution of Eq. (10) for $J_0 = 1$, $\alpha = 0.1$; the horizontal line denotes the critical value $A(T) = \lambda_{max} = J_0/\sqrt{\alpha}$.

The solution $A(T)$ has two branches $A^\pm(T)$ displayed in Figure 2. Since $A(T)$ is always larger than $J_0/\sqrt{\alpha}$ on the upper branch ($A^+(T)$), within this approximation the high temperature phase is *locally* stable down to zero temperature. It is therefore impossible to reach the low temperature phase from high temperatures without a discontinuity in $A(T)$. A lower limit on the discontinuity is the gap between two branches $\delta A = \frac{1}{2}\sqrt{\alpha}J_0 \ll J_0/\sqrt{\alpha}$ (recall that we are taking $\alpha \ll 1$). We note that although the feedback terms are generally small, they become important near the transition and here change its order. We can estimate the corresponding *minimal* jump in $|m_i|^2$

neglecting the feedback effects on the quartic terms (i.e. using $F^{(2)} = [A(T) - \lambda_{max}]|m|^2$, $F^{(4)} = \frac{1}{4}T|m_i|^4$), to be $\delta|m_i|^2 = \alpha$. In the vicinity of $T_{c0} = \lambda_{max}$ the discontinuity becomes larger ($\Delta A = \lambda_{max} - A^-(T_{c0}) = J_0/\sqrt{2}$); since we expect the transition to occur in the vicinity of T_{c0} , the most likely value of magnetization jump is $\Delta|m_i|^2 = \sqrt{2}\alpha$, as is also borne out by a more sophisticated calculation [10].

In the regime of strong frustration, when $\alpha \sim 1$, the transition becomes strongly first order. The case $\alpha = 1$ has a special interest because in this situation $\hat{J}$ becomes a unitary matrix and cumulants of $\hat{J}$ resemble those of the SK model: $\frac{1}{2N}\text{Tr}\hat{J}^{2n} = J_0^{2n}$. As a result, the calculation becomes simpler. The leading order locator equation (10) can be solved explicitly

$$A_{\alpha=1}^\pm = \frac{1}{2}[T \pm \sqrt{T^2 + 4J_0^2}] \quad (11)$$

suggestive of a strong first order transition around $T \sim J_0$. Certainly, the existence of such a transition cannot be determined by a m -expansion. Instead, we adopt a variational approach and demonstrate that there exists a “magnetic” phase whose free energy is lower than the “paramagnetic” free energy $F^+ = -2NTS_0(T)$ associated with the upper branch $A^+(T)$. We choose the trial state

$$m_j^{(v)} = \exp(-i\frac{\pi}{N}\gamma j^2) \quad (12)$$

$$m_k^{(h)} = (i)^{1/2} \exp(i\frac{\pi}{N}\gamma^{-1}k^2) \quad (13)$$

where γ is a variational parameter; since $|m_j|^2 = 1$ the free energy of this state is equal to its energy

$$\begin{aligned} E &= -J_0 \sum_{j,k} \frac{(-i)^{1/2}}{\sqrt{N}} \exp[-i\frac{\pi}{N}(\gamma j - k/\gamma)^2] + c.c. \\ &= -2N\epsilon(\gamma)J_0 \end{aligned} \quad (14)$$

where the dimensionless function $\epsilon(\gamma)$ satisfies $\epsilon(\gamma) = \epsilon(1/\gamma)$ and $\epsilon(\gamma) = \gamma$ at $\gamma \ll 1$. Clearly at $T \ll J_0$ this state has a lower free energy than the paramagnetic state associated with $A^+(T)$, and thus we conclude that it is unstable at low temperatures.

We emphasize that it is difficult to observe a paramagnetic instability in the free energy expanded in m_i ; we have already seen that its quadratic part remains positive definite on the upper branch $A^+(T)$. Now we show that this conclusion does not change when quartic terms are taken into account. To do so we assume the most general form of the free energy possible at $\alpha = 1$

$$\begin{aligned} F\{m_j\} &= -\sum_{kj} m_k^* \mathcal{J}_{kj} m_j + \sum_j [A(T)|m_j|^2 + \frac{1}{4}b|m_j|^4] \\ &\quad - \sum_j (h_j^* m_j + c.c.) + \frac{1}{2N} \sum_{kj} a_{kj} |m_k|^2 |m_j|^2 \end{aligned} \quad (15)$$

where the coefficients $a_{kj} = a_+$ if $\{k, j\}$ label parallel wires and $a_{kj} = a_-$ otherwise. We solve the equation $\frac{\partial F\{m_j\}}{\partial m_k} = 0$ to find the nonlinear susceptibility $\chi_{kj}^{(3)} \equiv \frac{\partial^3 m_k}{\partial h_k \partial h_j \partial h_j^*}$; it also satisfies $\chi_{kj}^{(3)} = -\frac{1}{T} |\chi_{kj}|^2 + O(|m|^2)$. Comparing these two expressions we determine the coefficients a_+ , a_- and b :

$$\begin{aligned} b &= T \\ a_+ &= \frac{J_0^4}{A^4(T) - J_0^4} T \\ a_- &= -\frac{J_0^2 A^2(T)}{A^4(T) - J_0^4} T. \end{aligned} \quad (16)$$

The quartic term of the free energy is minimized when $|m_j|^2 = m^2$, but still remains positive for the upper branch ($A^+(T) > J_0$)

$$F(m) = 2N \left[(A(T) - J_0^2) m^2 + \frac{1}{4} T \frac{A^2(T) - J_0^2}{A^2(T) + J_0^2} m^4 \right] \quad (17)$$

implying that all solutions remain stable to quartic order.

In summary, we have presented a non-random model which exhibits a first-order transition to a phase where the number of metastable states is extensive, consistent with one's general expectations for glassiness in the absence of disorder [1]. In this array the interaction between wires is long-range, and thus the barriers separating the low-temperature metastable states scale with the size of the system. Since the number of nearest-neighbors z is large we could perform an initial investigation using a modified TAP approach; furthermore the glass transition is weakly first-order in the limit of small α ($\frac{1}{N} \ll \alpha \ll 1$), the flux per unit strip, so that it could be studied analytically.

The results presented here can be tested in numerical simulations. The system will have a signatory superfluid stiffness, ρ_s , which can be measured by applying a twist Φ to the the horizontal(vertical) wires at their respective boundaries [11]; we predict that $\rho_s = \frac{\partial^2 F}{\partial \Phi^2}$ will appear discontinuously at the transition $T_c(\alpha)$ and that the jump will scale with α . Thermal cycling should indicate an extensive number of metastable states, thereby confirming the low-temperature glassy phase predicted here.

This transition could also be probed in fabricated arrays with negligible induced fields; more specifically we require

$$J_0 < \frac{\Phi_0^2}{N^{5/2} l} \quad (18)$$

which constrains the size of the array [12]. Here $T_c(\alpha)$ should be observable by resistive measurements. Alternatively one can infer the superfluid stiffness from inductive experiments. The critical current below $T_c(\alpha)$ should jump as well.

Of course, a glass transition is best characterized by its dynamical features, e.g. relaxation spectrum, hysteresis, ageing and memory effects; we have not yet investigated these properties of the periodic array discussed here. Hopefully the order parameter associated with this glass transition will also emerge from such a study, since we expect it to be a dynamical quantity. The thermodynamic behavior of a similar array with a reduced number of neighbors is another open question which could be important in understanding the relevance of long-range models to experimental glasses.

We would like to acknowledge M.V. Feigel'man for several useful discussions. We also thank the University of Oxford, the Aspen Center for Physics and NEC Research Institute for their hospitality. Partial financial support at Oxford was provided by SERC UK under Grant GR/H6680.

-
- [1] For a review of recent progress see G. Parisi, cond-mat preprint 9412034.
 - [2] We also note that in contrast to conventional 2D Josephson arrays, vortices are not independent degrees of freedom in this network as they are linearly confined.
 - [3] V. M. Vinokur, L. B. Ioffe, A. I. Larkin and M. V. Feigelman, Sov. Phys. JETP **66**, 198 (1987); M. V. Feigelman and L. B. Ioffe, cond-mat preprint 9405060.
 - [4] Note that for $\alpha \lesssim 1/N$ ferromagnetic ordering results. For $\alpha = 0$ the system corresponds to the conventional infinite-range ferromagnet albeit that J_0 must itself acquire an extra $1/\sqrt{N}$ dependence to yield the usual N -scaling of physical observables.
 - [5] D. J. Thouless, P. W. Anderson and R. G. Palmer, Phil. Mag. **35**, 593 (1977).
 - [6] We checked numerically that these results are correct for finite-size systems ($N \sim 50$) as well.
 - [7] We have not explicitly calculated all the states beneath the transition but it seems highly probable that the degeneracy of the soft mode will lead to a corresponding degeneracy of states.
 - [8] For a review of Potts spin glasses which exhibit such jumps see D. Sherrington, Prog. Theor. Phys. (Japan) Suppl. **87**, 180 (1986).
 - [9] M. V. Feigelman and A. M. Tselik, Sov. Phys. JETP **50**, 1222 (1979); A. J. Bray and M. A. Moore, J. Phys. **C12**, 441 (1979).
 - [10] To derive the discontinuity in $|m_i|^2$ we evaluate the free energy $F\{m_i\}$ to all orders in $q = \frac{1}{2N} \sum |m_i|^2$ but expand in $(|m_i|^2 - q)$. Finally, q itself is obtained by minimizing $F(q)$.
 - [11] S. Teitel and C. Jayaprakash, Phys. Rev. Lett. **51**, 1999 (1983).
 - [12] This condition was not satisfied by the experiment described in L. L. Sohn, M. T. Tuominen, M. S. Rzchowski, J. U. Free and M. Tinkham, Phys. Rev. **B47**, 975 (1993).